



EURAP

An International Antiepileptic Drugs and Pregnancy Registry

Interim Report – November 2019

Central Study Coordinator

Dr. Dina Battino

Department of Neurophysiology and Experimental Epileptology

Fondazione IRCCS Istituto Neurologico Carlo Besta

*20 133 Milano, **Italy***

Tel: + 39-02-23-94-22-30

Fax: + 39-02-700-42-91-60

E-mail: eurap@istituto-besta.it

Chairman Central Project Commission

Dr. Torbjörn Tomson

Department of Neurology

Karolinska University Hospital

*S-171 76 Stockholm, **Sweden***

Tel: + 46-08-51-77-37-05

Fax: + 46-08-51-77-37-57

E-mail: torbjorn.tomson@sll.se

BACKGROUND

A number of independent groups with experience and interest in maternal and foetal well-being in association with maternal use of antiepileptic drugs (AEDs) have agreed on a prospective international multi-centre study of pregnancies with AEDs. Data from all participating groups are shared in a Central Registry of Antiepileptic Drugs and Pregnancy (EURAP). EURAP was established in the first centres in some European countries and has since then gradually expanded to include more centres and countries now involving also Asia, Oceania, Latin America and Africa.

OBJECTIVE OF EURAP

The primary objective of EURAP is to evaluate and determine the comparative risk of major foetal malformations following intake of AEDs and their combinations during pregnancy.

METHODS

EURAP is an observational study. Women taking AEDs at the time of conception, irrespective of the indication, may be included. To avoid selection bias, only pregnancies recorded before foetal outcome is known and within week 16 of gestation are included in the prospective risk assessment. Cases ascertained later in pregnancy are recorded as retrospective cases, as they may provide signals, but are not included in the comparative risk evaluation.

Information on patient's demographics, type of epilepsy, seizure frequency, family history of malformations, drug therapy and of other potential risk factors is obtained, and follow-up data are collected once at each trimester, at birth and at one year after delivery.

Networks of reporting physicians have been established in countries taking part in the collaboration. During the course of the pregnancy, and the follow-up time after delivery, the participating physician enters data into five Subforms (Subforms A-E) for each patient.

Subform A is completed on enrolment of the patient, Subform B after the first trimester, Subform C after the second trimester, Subform D within three months after delivery, and Subform E within 14 months after birth. Immediately after completion, each Subform is submitted to the national coordinator for review. The national coordinator transfers the reviewed and accepted Subform to the Central EURAP Registry in Milan, Italy.


EVALUATION OF OUTCOME

The physician records descriptively abnormalities observed in the offspring. The final assessment and classification of the type of malformation is the responsibility of the Central Project Commission (CPC). In order to facilitate a uniform and objective assessment, reports of malformations are assessed regularly by an outcome assessment committee, which is kept blinded with respect to the type of exposure.

INTERIM REPORT

EURAP was implemented in the first two countries in Europe in 1999 and has since then grown to include countries from Europe, Oceania, Asia, Latin America and Africa. This development is reflected by increasing numbers of enrolled pregnancies. The development since 1999 is illustrated in Fig. 1.

Fig 1. Number of Participating Countries and Pregnancies Reported to the Central Registry by September, 2019.



The present report is **based on data available in the Central Registry by December, 10th, 2019**. At that time more than 1,500 reporting physicians from 45 countries had contributed cases to the Central Registry. Countries that have contributed at least 10 pregnancies in the current report are listed in Table 1.

Table 1. Countries that have contributed at least 10 pregnancies in the current report (n=34).

COUNTRY	National Coordinator (or referring physician*)	Date of joining the Registry
Argentina	Silvia Kochen	2002
Australia	Frank Vajda	2000
Austria	Gerhard Luef	2000
Belgium	Dick Lindhout & Eugène van Puijenbroek	2002
Chile	Alejandro De Marinis	2002
China	Weiping Liao	2006
Croatia	Dinko Vitezic	2002
Czech Republic	Jana Zarubova	2001
Denmark	Anne Sabers	2000
Finland	Reetta Kälviäinen	2003
France	Aileen McGonigal*	2000
Georgia	Sofia Kasradze; Nino Gogarishvili*	2000
Germany	Bettina Schmitz	2000
Hong-kong	Patrick Kwan	2002
India	Sanjeev Thomas	2001
Israel	Miri Neufeld	2000
Italy	Luigi M. Specchio	2000
Japan	Hideyuki Ohtani	2001
Lithuania	Ruta Mameniskiene	2002
Macedonia	Gordana Kiteva Trenevska	2001
Netherlands	Dick Lindhout & Eugène van Puijenbroek	2002
Norway	Silje Alvestad	2000
Philippines	Leonor Cabral-Lim	2003
Poland	Joanna Jedrzejczak	2001
Portugal	Isabel Pires*; Joana Parra*; Ines Cunha*; Elia Baeta*; Carla Bentes*; Catarina Cruto*	2001
Serbia & Montenegro	Maja Milovanovic	2002
Slovakia	Vladimír Safcák	2002
Slovenia	Boštjan Čebular & Gal Granda	2002
Spain	Meritxell Martinez Ferri	2001
Sweden	Torbjörn Tomson	2000
Switzerland	Barbara Tettenborn & Martin Kurthen; Dominique Flügel*	2001
Taiwan	Hsiang-Yu Yu	2004
Turkey	Demet Ilhan Algin	2000
United Kingdom	John Craig & Craig Heath	2001

* referring physicians

By the cut-off date for this report (December, 10th, 2019), **26,136 pregnancies had been entered into the central database**. Of these, **11,256 pregnancies are excluded** from the present interim report for reasons explained here below:

1. Pregnancies that failed to meet inclusion criteria (n= 193).
2. Lost to follow-up, including those failing to submit sub-forms within preset deadlines (n= 3,368).
3. Pending pregnancies, awaiting updates or corrections of different sub-forms (n= 1,204).
4. Ongoing pregnancies, updated and corrected (n= 529).
5. Retrospective, but completed and corrected (n=4,342). Among these, there are true retrospective pregnancies (n=4,019) and a further three hundred and twenty three pregnancies (n=323) that otherwise met our criteria for prospective pregnancies since they were recruited within 16th week, but for which patients had an ultrasound examination performed before enrolment.
6. Retrospective, i.e. initially classified as prospective pregnancies but re-classified as retrospective cases because one or more CRF subforms were submitted after the set deadlines (n=394).
7. Unclassifiable i.e. cases for which it was impossible to determine if there was a malformation or not (n=83). This includes 1 stillbirth with unknown fetal status, induced abortions with insufficient information on fetus (n=6), anomalies in livebirths where the information was insufficient to determine if qualifying for malformation diagnosis (n=71), 1 incomplete spontaneous abortion with unclear results of biopsy, and 4 perinatal deaths in premature births (<35 gestational weeks) with anomalies difficult to classify as congenital or due to prematurity.
8. Not yet classified, i.e. pregnancies which classification is pending as well as pregnancies which became completed after the last time we sent the database to the Outcome Assessment Committee (OAC), regardless if they contained some malformations or not (n=76).
9. Treatment changes between different AEDs or mono- to polytherapy or vice versa during the first trimester (n=1,067).

Thus in total **14,880 prospective pregnancies** (enrolled at the latest during the 16th gestational week and before outcome is known) **are included** in this report.

The classification of the epilepsy among the prospective pregnancies is given in table 2. Epilepsy was the indication for treatment in all but 112 (1%) of the pregnant women.

Table 2. Classification of the epilepsy in 14,880 prospective pregnancies.

Epilepsy	N	%
Localisation-related*	7,835	52.6
Generalized	6,162	41.4
Undetermined	486	3.3
Missing information	285	1.9
No epilepsy	112	0.8
Total	14,880	100

**Focal, according to more current terminology.*

The **maternal age** among prospective cases was **30.1 ±5.1 years** (mean±SD), ranging from 14 to 55 years.

The women were of Caucasian **ethnicity** in 87% and of Asian in 10%.


Gravida for each pregnancy is presented in Table 3.

Table 3. Number of the pregnancy in prospective cases.

Gravida	N	%
1st pregnancy	6,782	45.6
2nd pregnancy	4,645	31.2
3rd pregnancy	2,053	13.8
4th pregnancy	860	5.8
5th pregnancy	324	2.2
> 5th pregnancy	213	1.4
Not ascertained	3	0.0
Total	14,880	100

The outcome of the prospective completed pregnancies is presented in Figure 2. Out of the **277 induced abortions**, 43 were for chromosomal abnormalities and/or syndromes and 71 were for other fetal indication detected by prenatal screening (*out of these 71 cases, 59 were confirmed as major malformations and the remaining 12 cases were definitively classified as other abnormalities such as hydrops fetalis, molar pregnancies, blighted ovum, fetal placental transfusion syndromes, fetal growth retardation, fetus papyraceus, fetal death for unspecified causes, balanced translocation and insertion in normal individual*).


Figure 2. Obstetrical outcome of prospective pregnancies.



Of the pregnancies, **11,905 (80%) involved women on a single AED**, 2,387 (16%) were on two AEDs whereas 421 (2.8%) took three AEDs or more. One hundred and sixty-seven women (1.1%) were not on AED treatment during the 1st trimester. The exposure to the different AEDs in monotherapy among the prospective pregnancies is presented in Figure 3.

Figure 3. Number of prospective pregnancies with exposure to different AEDs in monotherapy.

Monotherapies (n=11,905)



There were 316 different AED combinations. The most frequently used combinations were lamotrigine and levetiracetam (n=311), lamotrigine and valproic acid (n=282), carbamazepine and levetiracetam (n=158), carbamazepine and lamotrigine (n=122), carbamazepine and clobazam (n=116), lamotrigine and topiramate (n=95), carbamazepine and valproic acid (n=83), carbamazepine and phenobarbital (n=82), clobazam and lamotrigine (n=59), levetiracetam and oxcarbazepine (n=59), clonazepam and lamotrigine (n=56), levetiracetam and valproic acid (n=54), and carbamazepine and topiramate (n=52) (Table 4).

Table 4. The most common AED combinations.

The most common polytherapies	N
lamotrigine + levetiracetam	311
lamotrigine + valproic acid	282
carbamazepine + levetiracetam	158
carbamazepine + lamotrigine	122
carbamazepine + clobazam	116
lamotrigine + topiramate	95
carbamazepine + valproic acid	83
carbamazepine + phenobarbital	82
clobazam + lamotrigine	59
levetiracetam + oxcarbazepine	59
clonazepam + lamotrigine	56
levetiracetam + valproic acid	54
carbamazepine + topiramate	52
lamotrigine + oxcarbazepine	41
clonazepam + valproic acid	39
topiramate + valproic acid	38
phenobarbital + valproic acid	38
levetiracetam + topiramate	32
carbamazepine + clonazepam	32
phenobarbital + phenytoin	32
clobazam + oxcarbazepine	30
lamotrigine + phenobarbital	27

The number of pregnancies with exposure to different second generation AEDs taken in combination with other AEDs are listed in Table 5.

Table 5. Number of pregnancies with different second generation AEDs in combination therapy.

Lamotrigine	1,287
Levetiracetam	896
Topiramate	372
Oxcarbazepine	251
Zonisamide	77
Gabapentin	64
Lacosamide	62
Vigabatrin	37
Pregabalin	28
Eslicarbazepine-acetate	14
Tiagabine	10
Perampanel	5
Rufinamide	2
Brivaracetam	2
Retigabine	1

TERATOGENIC OUTCOME

There were 673 major congenital malformations (MCM), 22 syndromic and/or genetic cases and 84 chromosomal abnormalities (CHR) in the prospective cohort of 14,033 pregnancies as shown in Table 6 (847 spontaneous abortions are excluded).

Table 6. Pathological outcomes.

Outcome	Outcome Classification	N
MCM	Multiple major	54
	Isolated major	619
MCM		673
SYNDROMES or GENETIC conditions		22
CHR		84
Total		779

The 22 syndromic and/or genetic cases are Marfan's syndrome (2), Noonan syndrome (2), inherited tuberous sclerosis (6), Goldenhar syndrome (1), incontinentia pigmenti (2), inherited congenital glaucoma (1), inherited congenital cataract (1), inherited craniosynostosis (1), Di George's syndrome (1), bilateral hearing loss (1), X-linked lissencephaly (1), Skeletal dysplasia/Dwarfism (1), X-linked ichthyosis (1) and Freeman Sheldon syndrome (1).

In this report we will confine our analysis to the 673 MCM including 59 induced abortions, six stillbirths and 16 neonatal deaths. Of the 592 live births, 75 cases of malformations were ascertained prenatally, 353 were first reported at birth, and a further 164 cases not detected at birth but within one year after birth.

Among the 673 cases with MCM, 143 were detected by ultrasound examination. Out of these 143, there were 59 induced abortions, five stillbirths, four perinatal deaths and 75 live births.

The 673 cases represent a [malformation prevalence of 4.8%](#) of all prospective pregnancies for which follow-up has been completed (673/14,033).

The type of malformations is described in Table 7.

Table 7

PATHOLOGICAL OUTCOMES	DESCRIPTION	N
MCM	Multiple major	54
	Nervous system	
MCM	Spina Bifida	42
MCM	Anencephalus and similar	4
MCM	Hydrocephaly	7
MCM	Microcephaly	2
MCM	Nervous system (other malformations)	15
	all	70
	Cardiovascular system	
MCM	Atrial septal defect	37
MCM	Ventricular septal defect	56
MCM	Atrioventricular septal defect	2
MCM	Congenital heart disease	51
MCM	Tetralogy of Fallot	4
MCM	Transposition of great vessels (complete)	4
MCM	Pulmonary valve stenosis	10
MCM	Hypoplastic left heart	8
	all	172
	Urinary system	
MCM	Urinary system (other malformations)	48
MCM	Renal Dysplasia	5
	all	53
	Digestive system	
MCM	Diaphragmatic hernia	8
MCM	Ano-rectal atresia and stenosis	2
MCM	Digestive system (other malformations)	10
MCM	Duodenal atresia or stenosis	2
MCM	Gastroschisis	3
MCM	Omphalocele	3
MCM	Atresia of oesophagus without fistula	1
	all	29
	Limbs	
MCM	Upper limb reduction	8
MCM	Lower limb reduction	1
MCM	Syndactyly	6
MCM	Polydactyly	24
MCM	Club foot - talipes equinovarus	19
	all	58
	Musculoskeletal	
MCM	Musculo-skeletal (other malformations)	11
MCM	Hip dislocation and/or dysplasia	70
	all	81
	Genital system	
MCM	Genital (developmental ovarian cyst)	6
MCM	Hypospadias	79
	all	85
	Eye, ear, face and neck	
MCM	Congenital cataract	4
MCM	Eye (other malformations)	3
MCM	Ear, face and neck	5
MCM	Choanal atresia	1
MCM	Atresia of nasopharynx	1
	all	14
	Oro facial clefts	
MCM	Cleft lip with or without palate	15
MCM	Cleft palate	16
	all	31
MCM	Other specified malformations (including sacral teratoma, cystic hygroma, haemangiomas, accessory skin tags, aberrant subclavian artery, congenital malformation of spleen, sequences, genetic syndromes, congenital malformation of renal artery, congenital malformation of adrenal gland, congenital malformations of integument, congenital malformations of lung)	26
MCM	all MCMs	673
	Chromosomal	
CHR	Chromosomal	22
CHR	Down's syndrome	41
CHR	Edward syndrome/trisomy 18	8
CHR	Klinefelter's syndrome	2
CHR	Patau syndrome/trisomy 13	6
CHR	Turner's syndrome	4
CHR	Wolff-Hirschorn syndrome	1
CHR	all CHR	84
	Syndromes or genetic conditions	
Syndrome	Marfan's syndrome	2
Syndrome	Incontinentia pigmenti	2
Syndrome	Noonan's syndrome	2
Syndrome	Goldenhar syndrome (Oculo-auriculo-vertebral syndrome)	1
Syndrome	Di George's syndrome	1
Syndrome	Tuberous sclerosis	6
Syndrome	Craniosynostosis, inherited	1
Syndrome	Congenital cataract, inherited	1
Syndrome	Congenital glaucoma, inherited	1
Syndrome	X-linked Ichthyosis	1
Syndrome	X-linked Lissencephaly	1
Syndrome	Hearing loss, bilateral, inherited	1
Syndrome	Skeletal dysplasia (achondroplastic Dwarfism)	1
Syndrome	Freeman Sheldon Syndrome (distal arthrogyrosis type 2A)	1
Syndromes	all Syndromes	22
Total	all cases with pathological outcomes	779

In 496 out of 11,266 pregnancies with AED monotherapy, one or more MCMs were observed (4.4%) as opposed to 172 out of 2,606 pregnancies with AED polytherapy (6.6 %), as shown in Table 8.

Table 8. Pathological outcomes by AED treatment categories.

(In this table, 847 spontaneous abortions have been excluded from the denominator).

	No AED	%	Monotherapy	%	Polytherapy	%	Total
MCM	5	3.1	496	4.4	172	6.6	673 (4.8%)
CHR	1	0.6	68	0.6	15	0.6	84 (0.6%)
Syndromes	0	0.0	17	0.2	5	0.2	22 (0.2%)
No malformation	155	96.3	10,685	94.8	2,414	92.6	13,254 (94.4%)
Total	161	100	11,266	100	2,606	100	14,033 (100%)

PUBLICATIONS

Changes in AED prescribing patterns and in rates of MCM over time in the EURAP cohort were published in *Neurology*. 2019 Aug 27;93(9):e831-e840.

Outcome regarding the eight most common monotherapies has been published in *Lancet Neurology*, April 18, 2018.

The dose-dependent risk of MCM with exposure to valproate in mono- and polytherapy has also been analysed and reported (*Neurology*, Sept 8, 2015) and so has the risk of intrauterine death in association with different treatments (*Neurology* Aug 18, 2015).

A manuscript on seizure control in pregnancies with withdrawal of or switch from valproate during 1st trimester as compared with maintained valproate treatment has been published in *Epilepsia* (*Epilepsia* 2016; 57: e173-7).

Outcome in relation to exposure to individual drugs or specific drug combinations is not included in the present report.

ORGANISATION, FUNDING AND SUPPORT

EURAP is a consortium of independent research groups working on a non-profit basis. The project is administratively organised by the Central Project Commission (CPC) with members representing different geographical areas and disciplines. The project has been supported by educational grants to the CPC from Eisai Pharmaceuticals, GlaxoSmithKline, Janssen-Cilag, Johnson & Johnson, Pfizer, Bial, Sanofi-Synthelabo, Novartis,UCB Pharma, and Teva. In addition, national and regional networks may receive support from the same or other pharmaceutical companies.

APPENDIX

Central Project Commission

Dina Battino, Milano

Erminio Bonizzoni, Pavia

John Craig, Belfast

Dick Lindhout, Utrecht

Emilio Perucca, Pavia

Anne Sabers, Copenhagen

Sanjeev V Thomas, Trivandrum

Torbjörn Tomson, Stockholm, (chair)

Frank Vajda, Melbourne

Central Study Coordinator

Dina Battino, Milan

Scientific Advisory Board

Bernd Schmidt, Freiburg

Martin J Brodie, Glasgow

Outcome Assessment Committee

(The persons below have contributed to the work of the OAC during different time periods of the project)

Chiara Pantaleoni, Milan, Italy

Elisabeth Robert-Gnansia, Lyon, France

Francesca Faravelli, Genoa, Italy

Richard Finnell, Houston, Texas